

Electric fish genomics: Progress, prospects, and new tools for neuroethology



William R. Pitchers*, Savvas J. Constantinou, Mauricio Losilla, Jason R. Gallant*

Dept. of Integrative Biology, Michigan State University, 288 Farm Lane RM 203, East Lansing, MI 48824, USA

ARTICLE INFO

Article history:

Received 1 July 2016

Received in revised form 6 September 2016

Accepted 16 October 2016

Available online 18 October 2016

Keywords:

Electric fish

Genomics

Transcriptomics

Genome editing

CRISPR

Transgenics

ABSTRACT

Electric fish have served as a model system in biology since the 18th century, providing deep insight into the nature of bioelectrogenesis, the molecular structure of the synapse, and brain circuitry underlying complex behavior. Neuroethologists have collected extensive phenotypic data that span biological levels of analysis from molecules to ecosystems. This phenotypic data, together with genomic resources obtained over the past decades, have motivated new and exciting hypotheses that position the weakly electric fish model to address fundamental 21st century biological questions. This review article considers the molecular data collected for weakly electric fish over the past three decades, and the insights that data of this nature has motivated. For readers relatively new to molecular genetics techniques, we also provide a table of terminology aimed at clarifying the numerous acronyms and techniques that accompany this field. Next, we pose a research agenda for expanding genomic resources for electric fish research over the next 10 years. We conclude by considering some of the exciting research prospects for neuroethology that electric fish genomics may offer over the coming decades, if the electric fish community is successful in these endeavors.

© 2016 Elsevier Ltd. All rights reserved.

Introduction

The powerful tools of genomics and genome manipulation are currently used nearly exclusively on a handful of well-established model species whose facile genetics and tractable husbandry allowed for the development of those tools. While this drives progress in those fields, the paucity of systems with reliable genomic data limits the insights that can be gained into the genetics of ecologically relevant traits, and may impede work on a number of central evolutionary problems. A core strength of the electric fish research, and of neuroethology as a discipline, is the focus on phenotype, broadly construed. This focus, combined with the strengths of electric fish as a model system, places neuroethology in a position to contribute to three of the five ‘Grand Challenges’ for biology as recently set out by the National Research Council (National Research Council, 2010) namely Connecting Genotype to Phenotype, Understanding the Brain, and Understanding Biological Diversity.

Since the discovery that weakly electric fish use electricity to sense their surroundings and communicate (Lissmann, 1958), researchers with interests and expertise spanning the range of bio-

logical disciplines have congregated around electric fish. Although the focus of contemporary work in the field has broadened considerably, electric fish researchers have continued to make valuable phenotypic and ecological connections that outpace many model organisms. The combination of genomics and the unique physiology of electric fishes – where the details of the electrosense link ecology and evolution intimately with neuroanatomy and ion channel kinetics (Fig. 1) – could allow for sweeping insights into how genotype connects to phenotype in an ecologically relevant system.

In the last few years, the availability of low-cost, high-throughput next-generation sequencing and sophisticated new molecular genetic techniques has laid the foundations for a ‘genomic renaissance’ in electric fish research. This paper will review the work of molecular biology in electric fish, beginning with the early biochemical contributions of *Torpedo* and *Electrophorus* and ending with whole genome sequencing efforts. In this work, the electric organ discharge (EOD) is a window into the neural system and the molecular workings of the electric organ (EO) and its component cells, and a view outward onto the ecology, behavior and evolution of the whole organism (Fig. 1). The latter part of the paper will then discuss the methods and benefits of integrating genomic and molecular tools into existing research programs.

* Corresponding authors.

E-mail addresses: pitchers@msu.edu (W.R. Pitchers), jgallant@msu.edu (J.R. Gallant).




Fig. 1. The 'integrated phenotype concept' in electric fish.

1. Molecular biology of electric fishes: the first 30 years

1.1. The molecular biology of the neuromuscular junction

Torpedo rays (Miledi et al., 1971; and to a lesser extent *Electrophorus electricus*; Changeux et al., 1970) contributed deeply to our understanding of synaptic transduction. With their large EOs – *Torpedo* is “essentially a swimming purified acetylcholine receptor” (Miller, 2000) – these species provide researchers with the abundant source of receptor-rich membranes needed for describing the structure and biophysical properties of proteins at the neuromuscular junction (NMJ). Given the connection between EO and motor plates (Keeseey, 2005), researchers could extract and purify the nicotinic acetylcholine receptor (nAChR) with relative ease (Sobel et al., 1977). These early biophysical studies were inherently comparative: immunohistological, ultrastructural (Rieger et al., 1976), and functional (Hess et al., 1982); however their efforts culminated in studies describing the detailed nanostructure of the nAChR (Kistler and Stroud, 1981), visualization of the molecular subunits forming the ‘rosette’ around the ionophoretic channel (Kistler et al., 1982), and finally a full 3D model of the molecule (Mitra et al., 1989). In addition, this research lead to the characterization of a number of other important proteins at the NMJ – notably agrin (Nitkin et al., 1987), dynein (Mou et al., 1998), and rapsyn (Elliott et al., 1980).

Beginning in 2007, Nazarian et al. (2011, 2007) demonstrated the potential for using genomic tools to probe the identity of NMJ proteins using high throughput Sanger sequencing. Nazarian et al. (2011) built a database of NMJ-associated proteins and transcripts present in electroplaques of *T. californica*, enabling the characterization of a suite of mammalian expressed sequence tags, and proteins of unknown function, as being associated with the NMJ. These discoveries proved crucial to the fields of biophysics and biomedicine (reviewed by Changeux (2012)), but although this work is indirectly relevant to neuroethology, it is the weakly electric fishes that have received the majority of the attention from the field.

1.2. Molecular biology of the EO: effectors and modulators

EOD duration varies 100-fold among species (Hopkins, 1999). This diversity is due in large part to variation in the electrocyte Na^+ currents (Ferrari et al., 1995), which in turn are regulated by voltage-gated Na^+ channels, and these are among the best studied electrocyte proteins. Research using polymerase chain reaction (PCR), quantitative PCR (qPCR), cloning, and first-generation sequencing has provided sequences of Na^+ channel genes from around 20 species of weakly electric fishes, representing both Gymnotiformes and Mormyroids (Arnégard et al., 2010; Lopreato

et al., 2001; Zakon et al., 2006). These studies provide a striking example of parallel molecular evolution: independent neofunctionalization (Ohno, 2013) of the voltage-gated Na^+ channel gene *scn4aa* in both lineages.

The teleost-specific whole-genome duplication event (Section 1.4) left these taxa with two paralogs (*scn4aa* and *scn4ab*) of the tetrapod muscle Na^+ channel gene $\text{Na}_v1.4$ (Lopreato et al., 2001; Novak et al., 2006). Following down-regulation of *scn4aa* in the ancestors of both lineages (Thompson et al., 2014), the expression of this paralog was found to be restricted to the EO (Zakon et al., 2006). Freed from the selective constraint to maintain the muscle phenotype (as *scn4ab* retained this function), *scn4aa* experienced strong positive selection on amino acid replacements critical for Na^+ channel kinetics, allowing for the evolution of EOD waveform diversification and signal complexity in both lineages (Arnégard et al., 2010; Zakon et al., 2008).

This subfunctionalization (evolutionary repurposing of duplicate genes; Magadum et al., 2013; Ohno, 2013) of membrane ion channels allows for greater EOD complexity and faster firing rates, but these properties are not fixed, even within a species. The EOD frequently varies between sexes and seasons, and is regulated by hormones (Bass and Hopkins, 1984; Hopkins, 1972). Furthermore, many studies have demonstrated that a variety of hormones have effects on the EOD (Allee et al., 2008; Bastian et al., 2001; Deemyad et al., 2013; Dulka et al., 1995; Dunlap et al., 2006; Dunlap and Zakon, 1998; Maler and Ellis, 1987; Markham et al., 2009a; Mills and Zakon, 1987; Smith and Combs, 2008; Telgkamp et al., 2007; Zupanc, 2002).

Changes to the EOD mediated by hormones have been recorded over a period of minutes (Markham and Stoddard, 2005), days or weeks (Dunlap et al., 1997; Ferrari et al., 1995; McAnelly and Zakon, 2007), and each is regulated differently. Changes occurring on the timescale of minutes are regulated by trafficking Na^+ channels into the electrocyte membrane (Markham et al., 2009b), and is circadian and socially controlled in *Sternopygus macrurus*. Circadian variation in EODs, at least in *Brachyhyppopomus* spp. (Franchina and Stoddard, 1998; Stoddard et al., 2007), is probably mediated by glutamate (Silva et al., 2008) and vasotocin (Perrone, 2010). Short-term hormonal EOD modulation seems to be mediated through cAMP and PKA after G-coupled receptor activation (Markham and Stoddard, 2005; McAnelly et al., 2003; McAnelly and Zakon, 1996).

Sexual dimorphism in EODs occurs in both Gymnotiformes (Allee et al., 2009; Ho et al., 2010; Markham and Stoddard, 2013; Smith, 2013) and mormyrids (Bass and Hopkins, 1985), and while androgens appear to be involved, other factors are implicated in mediating the dimorphism (Allee et al., 2009). Sexual dimorphism appears to have diverged rapidly, at least among *Apteronotus* spp. (Ho et al., 2010). The EOD (Carlson et al., 2000; Ho et al., 2010)

and its dimorphism (Franchina et al., 2001; Terleph, 2003) are also plastic in response to social cues, and this too appears to be hormone-mediated, with cortisol implicated (at least in *Brachyhyopomus*; Salazar and Stoddard, 2009). We refer readers to the excellent review by Markham (2013) for a more detailed exploration of this topic.

1.3. Evolution and development of sensory systems

Understanding electrolocation and electrocommunication requires studying not only EOs, but also the electroreceptive organs and the nervous system that serves them. To this end, numerous studies have used *in situ* hybridization or immunolabelling to localize a range of crucial molecules, including neurotransmitters and neurotransmitter receptors (Bottai et al., 1997; Johnston et al., 1990; Maler and Monaghan, 1991; Sas et al., 1990; Stroh and Zupanc, 1995); ion channels (Engelmann et al., 2008; Rashid et al., 2001); Ca^{2+} binding proteins (Friedman and Kawasaki, 1997; Smith et al., 2000) and neuropeptides (Sas et al., 1990; Stroh and Zupanc, 1995; Yamamoto et al., 1992; Zupanc et al., 1991). Zupanc (2002) reports more than fifteen neuroactive substances in just the central posterior/pre-pacemaker nucleus (CP/PPn) of gymnotiform fish of the genera *Apteronotus* and *Eigenmannia*. The central nervous system transcriptome of *Apteronotus leptorhynchus* has been assembled and validated by shotgun proteomics (Salisbury et al., 2015), thus producing a resource that may aid in exploring the underlying cellular mechanisms of critical neurobiological processes, such as adult neurogenesis, neuronal regeneration, and the neural basis of behavior. The origin of brain structures devoted to processing electrosensory information currently remain unknown (Carlson et al., 2011).

Electrosensitivity requires specialized sensory cells, and weakly electric fish have both high and low frequency tuned receptors (Gibbs, 2004; Jørgensen, 2005). In addition to bearing multiple types of receptors as an adult, there is evidence that the form and specificity of electroreceptors vary over an individual fish's growth and metamorphosis (Bensouilah et al., 2002; Denizot et al., 2007; Paintner and Kramer, 2003; Szabo, 1965). Given the apparent complexity of these systems, there is surprisingly little molecular data on electric fish electroreceptor development or the mechanism of electroreceptor signal transduction (unknown at the time of writing). Some expression data for RNA and protein in the electroreceptors of the weakly electric skate *Leucoraja erinacea* has been collected (Baker et al., 2013; Gillis et al., 2012; Maxwell et al., 2008), but there is more to be learned with transcriptomic/molecular techniques.

1.4. Evolution and development of novel electrogenic systems

Most EOs develop from myoblasts, i.e., skeletal muscle precursors (Kirschbaum, 1977; Szabo, 1960). As such, electrocytes share similarities with myocytes: they are multinucleated, excitable, and retain several muscular proteins. However, electrocytes are larger, non-contractile, more highly-polarized and more variable morphologically. How EOs have evolved from, and differ molecularly from skeletal muscle is a central question for electric fish researchers. EOs evolved independently in mormyrids and Gymnotiformes (Fig. 2) following the teleost-specific whole genome duplication event (Amores et al., 1998; Hoegg et al., 2004; Hurley et al., 2007; Loproto et al., 2001; Taylor et al., 2001). Studying the expression of the resulting paralogs across a range of species (e.g., Thompson et al., 2014) addresses the importance of sub-/neo-functionalization as a source of evolutionary novelty (Magadum et al., 2013; Ohno, 2013; Zhang, 2003).

The two best-studied aspects of EO differentiation are the dependence on innervation and the suppression of muscle-

specific protein expression. *Sternopygus macrurus* requires EO innervation for its differentiation and maintenance; denervating the EO causes the appearance of sarcomeric proteins in mature electrocytes, and inhibits differentiation of new electrocytes from satellite cells (Patterson and Zakon, 1993; Unguez and Zakon, 2002, 1998a; Weber et al., 2012; Zakon and Unguez, 1999). By contrast, the electrocytes of the Mormyrid *Pollimyrus isidori* do not require innervation for differentiation (Szabo and Kirschbaum, 1983), and histological observations on eight Gymnotiformes show that electrocytes differentiate before they are innervated (Kirschbaum and Schwassmann, 2008; Schwassmann et al., 2014).

Researchers have documented the mechanism of muscular protein suppression in *S. macrurus*. Initial immunolabelling and ultrastructural studies demonstrated that electrocytes express several muscle proteins, but lack some sarcomeric proteins, sarcomeres and T-tubules (Cuellar et al., 2006; Kim et al., 2004, 2008; Patterson and Zakon, 1996; Unguez and Zakon, 1998a, 1998b). Nevertheless, electrocytes transcribe the genes for the muscular proteins that they lack (Cuellar et al., 2006). Moreover, the myogenic regulatory factor genes, which control skeletal muscle gene expression, are all transcribed in the EO (Kim et al., 2008, 2004). The electrocyte transcript and protein profiles do not match, which suggests post-transcriptional mechanisms are responsible for the EO differentiation from muscle in *S. macrurus*.

Pinch et al. (2016) used genomic tools (both qRT-PCR and RNA-seq) to examine the regulation of sarcomere gene expression in *S. macrurus* by comparing EO and muscle transcription profiles of mRNA and miRNA. They did not detect differential expression in sarcomere-related genes between the two tissues, confirming earlier work in this species. In addition, protein-degradation pathways showed similar expression in both tissues, which implies that post-translational sarcomeric protein degradation does not regulate the electrocyte phenotype. They also identified 155 annotated miRNAs, all of which were expressed in both tissues. Three miRNAs were upregulated in the EO, and they were known to participate in muscle differentiation and the down-regulation of sarcomere-associated genes. The research on the non-contractile phenotype of electrocytes in *S. macrurus* unambiguously points toward a post-transcriptional mechanism of regulation via silencing a few sarcomeric genes.

While most data on EO developmental mechanisms has been collected in *S. macrurus*, these mechanisms may not be representative of all electric fish. For instance, *Brienomyrus brachyistius* expresses both transcripts and proteins from some muscle-specific genes while decreasing the expression of others (Gallant et al., 2012, 2014), suggesting that a transcriptional regulatory mechanism downregulates the expression of some muscle genes. Work on the mormyrids *Campylomormyrops compressirostris* and *Campylomormyrops tshokwe* (Lamanna et al., 2015), on the Gymnotiformes *E. electricus* and *Eigenmannia virescens*, and in the Siluriform *Malapterurus electricus* (Gallant et al., 2014), suggests that they share a similar transcriptional regulatory mechanism to that of *B. brachyistius*.

1.5. Unraveling the phylogeny of weakly electric fish

Electric fish can be problematic for taxonomists, with morphologically cryptic species not uncommon (Feulner et al., 2006; Nagamachi et al., 2010; Silva et al., 2015) and many phylogenetic details unresolved. To help resolve the relationships among species and populations, researchers have used genetic markers to infer relatedness. Microsatellites (or simple sequence repeats, SSRs) are short, repeated sequence motifs. These repetitions tend to accumulate mutations comparatively quickly and are therefore most used in situations where the relationships among subjects are expected to be close, such as assessing divergence among populations




Fig. 2. The distribution of available genomic data for electric fish. Genomes, transcriptomes, population genetics marker sets, partial sequences and proteomic datasets – ‘X’ indicates a single dataset, ‘XX’ indicates multiple datasets.

(Arnégard et al., 2005; Feulner et al., 2006, 2008). Amplified fragment length polymorphisms (AFLPs) are neutral markers used to detect polymorphisms among more distant relationships. DNA is fragmented using restriction enzymes, and the pattern of resulting fragments depends on the location of the restriction sites where the enzymes bind. Since these restriction sites evolve more slowly than SSRs, they are often used to infer relationships among species (Lavoué et al., 2008; Sullivan et al., 2004).

At larger scales, such as among genera, the sequence of the mitochondrial protein cytochrome *b* (Sullivan et al., 2004, 2000) and other mitochondrial sequences (Alves-Gomes and Hopkins, 1997; Lamanna et al., 2014; Lavoué et al., 2012) are used, since these parts of the genome evolve slower still. Molecular markers can also be used in combination with other potentially informative data such as the analysis of EOD recordings (Alves-Gomes and Hopkins, 1997; Feulner et al., 2008, 2006; Sullivan et al., 2000), geometric morphometrics (Feulner et al., 2008), and even fossil data (Lavoué et al., 2012).

Marker-assisted phylogenetic studies have sometimes supported existing species identifications based on morphology (Feulner et al., 2008; Sullivan et al., 2000) and have unmasked cryptic morphs elsewhere (Feulner et al., 2006). These studies also have the potential to reveal surprises, such as mitochondrial introgression between putative species (Lavoué et al., 2008; Sullivan et al., 2004) and a pattern of reduced variability among the EODs of Petrocephalines when compared to the Mormyrids. The potential for NGS technology to contribute to phylogenetic studies is demonstrated in a transcriptomic study where the complete mitochondrial genome for *C. compressirostris* was assembled and >1500 SSR markers were developed (Lamanna et al., 2014). With this density of markers, studies can effectively probe the structure of pop-

ulations and search for divergent regions or loci that show a signature of selection (Ellegren, 2014).

1.6. Macroevolution & diversification

The studies above demonstrate the integration between levels of analysis – molecular, cellular, and at the level of tissues/organs – that form the ‘integrated phenotype’ (Fig. 1). This integration is a key strength of weakly electric fish as a research system, but it can be extended; NGS techniques greatly facilitate the use of electric fish to address broad genetic and evolutionary questions. The ‘natural experiment’ in repeated independent evolution of the EO is well-suited for comparative molecular work, since the parallel evolution of Gymnotiformes and Mormyrids allows for the separation of phylogenetic ‘noise’ from the evolutionary ‘signal’.

Gallant et al. (2014) took this approach in examining the genomic basis of anatomical and physiological convergence across three independently evolved electric fish lineages. The authors assembled a draft version of the *E. electricus* genome and EO and skeletal muscle transcriptomes from three Gymnotiformes, one mormyrid and one Siluriform (see Section 1.4). Comparative analyses revealed genes potentially involved in the phenotypic characteristics of electrocytes that differentiate them from muscle, and suggested that a common regulatory network of transcription factors and developmental pathways may have been repeatedly utilized in the evolution of EOs (Gallant et al., 2014).

Work on the *Electrophorus electricus* genome (Gallant et al., 2014; Traeger et al., 2015) has also produced the first analysis of electric fish miRNA expression (Traeger et al., 2015). Researchers examined transcriptomes from eight adult tissues, including all three EOs and detected upregulated genes in these EOs that are

enriched in functions related to transmembrane transport, androgen binding, and receptor signaling. They also described a unique collection of expressed miRNA in each of the three EOs. This included three conserved miRNAs known to participate in the inhibition of mammalian muscle development, and one highly expressed novel miRNA.

2. Developing a ‘Genomic Toolkit’ for electric fish: a research agenda for the next decade

The integrated phenotype concept in electric fish (Fig. 1) makes them a powerful model system for addressing research challenges across biological disciplines, and more specifically in neuroethological contexts. Genomic and molecular data have accelerated in the past 5 years, largely due to the rise of low-cost next generation sequencing (NGS) approaches. The combination of the strengths of the electric fish model with coordinated, strategic advances in both data collection and protocol development, would greatly facilitate the ability to address 21st century research objectives (National Research Council, 2010), particularly with regard to understanding the brain, and integrating genotype with phenotype across levels of biological analysis. This coordinated research effort can be imagined as a ‘genomics toolkit’ consisting of established protocols to understand sequence divergence, assess and manipulate gene location/expression, and generate transgenic tools. The success of these endeavors over the next decade will undoubtedly usher in a new ‘golden age’ for electric fish biology. Such a genomic toolkit is only as useful as it is accessible, highlighting a need for community involvement and support for new standards for data sharing and protocol exchange.

2.1. Additional sequencing

The growth and coordination of sequence information are crucial for bringing electric fish biology into the new ‘golden age’ promised by genomic tools. At this point, genome sequencing should focus on species that can be bred in captivity (for production of transgenic lines), that are readily available, and that represent the principal taxonomic groups. Sequencing the genomes of fish with these characteristics will allow us to refine genetic tools, establish detailed protocols, and prime research for other electric fish species.

With NGS technologies becoming increasingly affordable, even small labs can acquire vast amounts of sequence data. Vertebrate genomes sequenced with 50× coverage can return 100s of gigabytes of data. Further, in the process of e.g., assembling a genome draft from this data, the space required for the various intermediate files can easily grow to more than a terabyte. Properly planning an NGS project therefore also involves planning for data management and backup. Many Universities maintain computing facilities and/or are connected to local or national grids, and we would encourage the use of such infrastructure where it is available. Where this isn't the case, an internet connection is all that is required to make use of cloud-based on-demand computing services, such as Amazon's [AWS](#) or [Rackspace](#). Sequencing centers themselves often offer some data management and bioinformatics support, but this varies dramatically among centers and so is a topic to discuss with your sequencing provider about while planning a project.

Sequencing transcriptomes may offer the best value for money in many situations, since transcriptomic data can be used to test for both expression and coding differences, in addition to identifying candidate loci for *in situ* hybridization and loss of function studies (see Sections 2.2–2.4). RNAseq may best be aimed at first generating nervous system expression profiles (e.g., of whole

brains, brain regions, peripheral neurons) for comparison to the *Apteronotus leptorhynchus* CNS transcriptome (Salisbury et al., 2015), and/or ontogenic expression profiles for developing tissues, e.g., electroreceptors. Whole genome sequencing would help identify targets for both loss of function studies and transgenic fishes, and comparative genomic studies could probe ‘big picture’ questions in the evolution of electrogenic fishes and, even more broadly, general biological processes.

A reduced representation sequencing technique, such as RAD-seq (for genomic DNA; Davey and Blaxter, 2011) or TagSeq (for RNA; Lohman et al., 2016) is comparatively inexpensive (compared to the cost of transcriptome or genome sequencing) for generating large sets of marker data with markers semi-randomly distributed throughout the genome, which is useful for phylogenetic studies.

Understanding this additional sequence data will require that researchers develop some new skills. While one does not necessarily need to be an expert bioinformatician to work with sequencing data, the vast majority of the software needed is run at the command line in a Unix environment, usually on a server or high-performance computing cluster. The development of genomics software moves fast, driven by the rapid pace of advance in sequencing technology – for that reason, any software recommendations here would very quickly be obsolete. That said, the ‘fastq’, ‘fasta’, ‘sam’ and ‘vcf’ file formats (see Table 1) are so widely used as to have become *de facto* standards, and researchers working with genomic data should familiarize themselves with these. What these formats have in common is that data is stored as plain text, files that are often too large to be opened on a personal computer. Working with data in these formats (and most other genomic formats) therefore principally involves programmatically manipulating files without opening them; for which the Unix environment is well-suited. Rather than recommending specific software packages, we would therefore encourage researchers to acquaint themselves with the Unix ‘[shell](#)’, and with at least one of the most commonly-used programming languages, such as [Python](#), [Perl](#) or [Ruby](#). For newcomers to genomics, a general computing skills text would be a good place to start (Haddock and Dunn, 2011). [Buffalo, 2015](#) are highly recommended), and a plurality of working bioinformaticians share tools and advice at websites like [Biostar](#), [SEQanswers](#) and [Stack Overflow](#).

2.2. Molecular biology applications post-sequencing

Sequence information has multiple uses. It can be used to generate PCR/qPCR primers for population genetic marker development (see Section 1.5), *in situ* hybridization templates, for differential expression and RNA profiling, comparative genomics, and identification of targets for loss of function studies and generation of transgenic lines.

Using specific gene sequence information, primers can be designed to amplify the focal gene specifically through PCR/qPCR. Identification of high quality PCR primers is necessary, due to the use for PCR in generating templates (e.g., for *in situ* probes) and assaying experimental outcomes. PCR can identify NGS-predicted alternative splice variants and be utilized to assay potential single nucleotide polymorphisms (SNPs). qPCR allows the quantification of DNA in real time showcasing its use as an assay of gene expression; however, qPCR can also be used for validating microarray experiments and monitoring biomarkers (VanGuilder et al., 2008).

With RNAseq, researchers can discover and quantify the full set of transcripts – including alternatively spliced isoforms and non-coding RNAs – from a sample (Martin and Wang, 2011). One common use for RNAseq data is differential expression analysis, wherein the expression level of every gene is compared statistically across two or more groups of samples (Mortazavi et al.,

Table 1

A brief glossary of genomics-related terms, and recommended reading for those wishing to learn more.

Genomics	
First-generation sequencing	'Sanger sequencing'; a long read technology still frequently used to sequence plasmids, PCR products and other relatively short DNAs
Next-generation sequencing (NGS)	A suite of technologies that use amplified sample DNA/RNA as a template (no need for sample cloning) and run millions of sequencing reactions in parallel. Sequence is produced in fragments
Library	A set of templates produced by shearing sample DNA into fragments and ligating oligo adapters at their free ends
Template	Recombinant DNA molecule made up of an adaptor sequence followed by the target sequence (typically an unknown fragment)
Adapters	Synthetic leading/trailing sequences added to fragments to facilitate sequencing chemistry
Barcodes	Synthetic sequences added to fragments (often as part of the adapter) to ID samples within a mixed-sample run
reads	The individual string of base-pair codes that are generated in parallel by NGS technologies. Require trimming and alignment before analysis
.fastq	The file format output by the vast majority of sequencing facilities. A plain text file that includes both base calls and a quality score for each base
.fasta	A plain text file format for containing named/labeled DNA or RNA sequences
Mapping/Alignment	The process of aligning short reads to a reference sequence, which could be a complete genome, transcriptome, or <i>de novo</i> assembly
<i>De novo</i> assembly	The computational rebuilding of the genome/transcriptome from the sequenced fragments
Assemblers	Software developed to tile sequenced fragments together – many tools available with various strengths/weaknesses
.sam/.bam	SAM is a format for storing sequence data in a series of tab delimited text columns. Currently the most common output format from mappers/aligners. BAM is a binary version: non-human-readable but much smaller files
.vcf	A plain text file format that lists only variant sites (rows) by a list of samples (columns)
annotation	Indexing of genomic and transcriptomic features by base-pair coordinates on raw genome or transcript sequence. Often formatted in .gff or .gtf format
transcriptome	Everything transcribed from the genome; primarily mRNA and rRNA, but containing many classes of small and large non-coding RNAs
Gene ontology	'GO' annotations: a way of assigning or analyzing functional gene annotations by comparing them to known biological pathways
'HiSeq' or 'miSeq'	Sequencing technologies by Illumina, and currently the standard at most sequencing facilities. Results in relatively short reads (up to 200 bp), with a lower error rate than PacBio. The principle advantage of these machines is their very high throughput; millions of reads per run
Ion Torrent	Sequencing technologies from ThermoFisher/Life Technologies. Similar in read length to Illumina (max. 400 bp), and often slightly more affordable, but is subject to increased errors in homopolymer runs. Yields many fewer reads per run and therefore comparatively low depth coverage
'PacBio'	Relatively new sequencing technology from Pacific Biosciences, that features very long reads (average 10,000 bp), but a much higher error rate than Illumina
RADseq	Restriction Associated DNAseq. By digesting genomic DNA with a specific enzyme before sequencing, a subset of the genome is sequenced with higher coverage. Useful for ascertaining intraspecies variation
TagSeq	The RNAseq technique of sequencing only the 3' end of mRNAs, such that the overall amount of sequence per individual is reduced, but the number of fragments sequenced is maintained. However, the ability to detect splice variants and some other features is lost
<i>Recommended reading</i>	
Bild et al., 2014; Borgman, 2012; Bradnam et al., 2013; Davey and Blaxter, 2011; De Wit et al., 2015; Earl et al., 2011; Ellegren, 2014; Ekblom and Galindo, 2010; Ekblom and Wolf, 2014; Field et al., 2009; Fonseca et al., 2012; Hatem et al., 2013; Kaye et al., 2009; Lee et al., 2016; Liu, 2005; Lohman et al., 2016; Macaulay and Voet, 2014; Martin and Jiggins, 2013; Metzker, 2009; Narum et al., 2013; Sims et al., 2014; Wake, 2008; Whitlock et al., 2010	
Wet-Lab Molecular Biology	
Genetic markers	Microsatellites, Simple sequence repeats, or AFLPs that can be easily assayed and used as a proxy or tag for the genotype of regions surrounding it
Neutral markers	DNA markers in non-coding and presumed non-regulatory regions
Comparative genomics	Comparing the structure, organization and gene content of genomes in a tree-wise fashion to elucidate trends and patterns of genome evolution
Splice variants	Multiple mature mRNAs derived from the same precursor RNA
miRNA	microRNA: a very short RNA molecule (~22 bp) involved in RNA silencing
RNAseq	Next-gen sequencing where RNA (transcriptome) is sequenced rather than DNA (genome). May be further refined to target specific families and types of RNAs
<i>In situ</i> hybridization	Allowing labeled DNA or RNA probes to anneal to DNA/RNA targets <i>in vivo</i> . Frequently used to localize mRNA transcripts to tissues
antisense	Complementary to an mRNA sequence. Double stranded mRNA is not effectively translated, and is targeted by the cell for degradation
knockdown	Targeting and downregulating or degrading a specific mRNA of a gene of interest. More attenuated effect than a knockout, which removes the gene entirely; avoids lethality and is easier to do than gene knockout
reporter	An attribute added to a mutant to access function or success of the mutation, e.g., adding GFP to a gene so its protein product can be localized by microscopy
<i>Recommended reading</i>	
Deisseroth, 2010; Deisseroth et al., 2006; Heidenreich and Zhang, 2015; Mancuso et al., 2010	

2008). This allows for the association of the differentially expressed genes with the differences between any two groups; e.g., different tissues, species, developmental stages, diseased or transgenic individuals, cell activation states. However, the identification of splice variants may also prove useful to neuroethologists in light of the emerging role of alternative splicing in neural patterning (reviewed by Li et al. (2007)).

The availability of genomic sequences for electric fish allows for the use of comparative genomics analyses. For instance, regions of

the genome that are highly conserved between divergent populations likely correlate with shared phenotypes. Selective events leave molecular signatures, (e.g., regions of reduced genetic diversity) that can be used to identify genes controlling EOD differences or other traits.

Further to these uses, sequence information is a prerequisite for planning follow-up molecular studies, such as loss of function manipulations (Section 2.3) and generating transgenic animals (Section 2.4).

2.3. Develop resources for transient loss of function studies

Loss of function studies enable researchers to assign a potential function to a gene by assaying how the cell/tissue/organism responds in the genes' absence. Transient loss of function studies mediate gene knockdowns via exogenously injected oligomers, allowing for transient effects on gene function. Tracking other gene/protein expression changes after transient loss of function studies can determine gene regulatory networks, highlighting again the remarkable capacity for large-scale biological processes (e.g., communication, sexual selection, neural coding) to be understood at the gene level in electric fish (Fig. 1). The most widely used oligomers that are injected to generate loss of function are morpholinos and double stranded RNA (dsRNA).

Phosphorodiamidate morpholino oligomers (or simply 'morpholinos') are small (~25 bp) oligomers that bind to specific RNA sequences, thereby interrupting post-transcriptional processing using an antisense technology (see review by Moulton (2016)). Knockdown of the targeted gene product can be achieved by blocking the translation of the targeted RNA sequence or by blocking the spliceosome and causing intron retention (Fig. 3). Morpholinos can be injected into single cell/developing embryos or adult tissues, without permanent effect. 'Scrambled' morpholinos are available to purchase for use as a control for morpholino-specific effects.

Vivo-morpholinos are synthesized with a carrier subunit that allows for delivery to target cells after intramuscular, intravascular, or intraperitoneal injections (Kotb et al., 2016; Notch et al., 2011; Schulman et al., 2016) and even into developing fish embryos with a bath immersion (Wong and Zohar, 2015). Modifications to the morpholino subunits can generate a tool for inducible gene knockdowns (Shestopalov et al., 2012), providing extra specificity and control; e.g., fluorescein-tagged morpholinos can be visualized as well as electroporated (Hyde et al., 2012). Correctly identifying targets for morpholino loss of function requires access to sequence data; either the coding sequence (translation-blocking morpholinos) or an annotated genome (to identify splice-blocking targets).

Like morpholinos, RNA interference (RNAi) is a type of antisense technology used to knockdown gene expression. RNAi consists of

injecting long double stranded RNA (dsRNA) or short dsRNAs (siRNAs, miRNAs, piRNAs) into an organism/cell. When the RNAi pathway is activated (Fig. 3), the dsRNA is cut by the enzyme DICER into small pieces, essentially generating many siRNAs that target the entire dsRNA sequence after forming the RNA-induced silencing complex (RISC) with other proteins (Alagia and Eritja, 2016; Kim et al., 2009). RNAi components have been manufactured that permit more precise spatiotemporal control and have been utilized in zebrafish (Andrews et al., 2014; Tang et al., 2007).

RNAi loss of function is similar to that of morpholinos; the effects are transient and affect protein expression through RNA, and thus could be used on species that only have transcriptomic data available. RNAi may be preferable to morpholinos due to lower costs and may be more viable for larger screens; e.g., De Rienzo et al. (2012) championed their use in zebrafish as being rapid in effect, inexpensive and a tissue-specific method beyond embryonic stages. However, RNA can be difficult to work with (as it is less stable than DNA) and cloning the sequence of interest is required.

One of the strengths of transient loss of function is the ability to induce loss of function in adult organisms. Because many electric fishes are difficult to breed in captivity, morpholinos and RNAi will likely be popular tools – and could easily be utilized in the field. These tools could be applied to understand how gene regulated changes to the EOD affect sexual selection, electrolocation, and electrocommunication.

2.4. Develop resources for manipulating the genome

While some studies require transient/post-developmental gene manipulation, other studies depend on constitutive gene expression or mutant lines. CRISPR/Cas9 (or simply CRISPR) enables researchers to generate transgenic lines and mutants. CRISPR originated as a prokaryotic immune system complex used as defense from foreign DNA (Barrangou et al., 2007), but has been modified for laboratory use. A single guide RNA (sgRNA) recognizes and binds with a specific sequence on the DNA along with a Cas9 endonuclease. This complex, through Cas9, then introduces double




Fig. 3. Schematic summary of morpholino, RNAi, and CRISPR approaches for gene/genome function manipulation. See text (Sections 2.3 and 2.4) for a description of these approaches. * Knockdown utilizing morpholinos and RNAi can also occur in adult tissues overtime (Section 2.3). ** CRISPR can also be performed via *in vitro* transcription of just sgRNA and subsequent injection of the sgRNA/Cas9 enzyme complex.

stranded breaks at that precise location. The double stranded break is repaired through the nonhomologous end-joining (NHEJ) pathway, ultimately rejoining the DNA but often introducing insertions, deletions, or both (indels), or, if a homologous donor sequence is present at the time of repair, the homology-directed repair (HDR) pathway can be activated and the genome may insert the donor sequence (Fig. 3; see review by Sander and Joung (2014)). Through NHEJ repair, mutant lines (knockouts) can be generated and HDR allows for transgenic lines to be produced.

One strength of CRISPR is that, unlike mutagenesis screens, specific genome regions can be targeted, and a range of knock-out/knock-ins can be made by modifying the genomic sequence *in vivo* (Mei et al., 2016; Ran et al., 2013; Tsai and Joung, 2016). Genomic edits are passed on to all the daughters of the treated cells; if incorporated into the gametes, genomic edits can be transmitted to all progeny, producing breeding lines with stable knock-out and knock-ins (Auer et al., 2014; Hwang et al., 2013; Kimura et al., 2014; Li et al., 2015). Knockouts through CRISPR can be highly efficient (75–99%) even for multiple genes in a single embryo (Jao et al., 2013). The effects are often biallelic, disrupting function completely and almost immediately (necessary for developmental studies). Current advances in CRISPR protocols have generated an inexpensive, cloning-free method for generating sgRNA and Cas9 mRNA/protein that most labs can readily adopt (Gagnon et al., 2014; see supplement for detailed protocol). CRISPR modifications occur in the genome, thus genomic sequence data is strongly recommended for identifying high quality CRISPR targets.

The potential applications for CRISPR have been expanded by recent molecular innovations (Harrison et al., 2014; Hsu et al., 2014). For example, using CRISPR it could be possible to visualize protein expression *in vivo* by inserting reporters, offering more information in concert with gene expression visualized by *in situ* hybridization. Green fluorescent protein (GFP) tagged proteins, coexpression of GFP with a gene of interest, or GFP under control of an endogenous gene promoter can be visualized with a fluorescent microscope to determine location of the gene of protein/transcript. Comparing both gene transcript and protein location can uncover surprising results (see Section 1.4); localizing both gene outputs in a single tissue would be much easier, and more informative, by performing *in situ* hybridization on a transgenic fish that always expresses fluorescent tagged protein for the gene of interest.

Not only can CRISPR facilitate protein localization through reporter knock-ins, but it can also be used to identify the sequence basis of protein properties. Zakon et al. (2008) note that we currently do not know how the positively selected substitutions on the electrocyte-expressed *scn4aa* alter the biophysical properties of the Na⁺ currents. CRISPR site-directed mutagenesis could identify the effect of single nucleotide polymorphisms on ion channel kinetics: either *in vivo*, or *in vitro* through exogenous expression (e.g., in a *Xenopus* oocyte). Such studies may have direct consequences beyond neuroethological and electric fish research, because several of these amino acid replacements occur near sites associated with disease-causing mutations in human Na⁺ channel genes (Zakon et al., 2006).

Similarly, CRISPR can replace not only SNPs, but entire gene sequences. Candidate genes for species divergence could be reciprocally inserted for definitive gene function assays, or to test for genetic effects of different genetic backgrounds. Linking changes in protein sequence to EOD signaling changes could facilitate studies exploring sexual selection and EO evolution among species; e.g., by driving male-like expression of candidate genes in female fish and *vice versa*. Finally, sgRNA's can target nearly any location in the genome, opening the possibility for studies on differences in non-coding/cis-regulatory regions.

2.5. Limitations of morpholinos, RNAi, and CRISPR

Transient loss of function tools and genomic editing are powerful; however, like any tool, each has its own limitations. Below we present several limitations, but researchers are encouraged to read the current literature on these issues while deciding which tool(s) to use.

Reports that compare morpholino efficiency to CRISPR or mutant strains (Kok et al., 2015) suggest there are potential off-target effects and phenotypes when using morpholino (but see Morcos et al., 2015). Like morpholinos, there are known off-target effects seen with RNAi (Jackson et al., 2003; Nunes et al., 2013); therefore, proper controls and careful interpretation of results are absolutely necessary.

The availability of single cell embryos is a necessity for CRISPR. Electric fishes of various species have been bred successfully, including multiple mormyrids and Gymnotiformes; Kirschbaum and Schugardt (2002) provide a relatively comprehensive list of all laboratory bred electric fish. Breeding of these fishes is not without difficulty; most freshwater electric fish breed during the rainy season and it is necessary to simulate some of these conditions to induce gonadal recrudescence. At a minimum, the water conductivity must be dropped (Schugardt and Kirschbaum, 2004), although some Gymnotiform species seem to also require an increase in water depth (Kirschbaum and Schugardt, 2002). Some effort has been made to generate developmental series for electric fishes (Diedhiou et al., 2007), allowing more rigorous developmental hypotheses to be tested in a systematic way among labs.

CRISPR has historical issues of non-specificity and off-target effects (Zhang et al., 2015). These have been addressed in part through the double-nickase approach (Mei et al., 2016; Ran et al., 2013; Tsai and Joung, 2016); however, this approach limits available targets. The knockdown efficiency of CRISPR can vary among target regions (Doench et al., 2014; Hua Fu et al., 2014), with some target genes currently inaccessible to genomic edits. Without developing inducible CRISPR constructs, embryonic-lethal genes cannot be thoroughly studied with knockouts, but titrations of dsRNA or morpholino could be used in such cases.

2.6. Community dissemination

Developing genomic tools for electric fish benefits the electric fish community as a whole and requires collaboration among researchers. Compiling and curating available electric fish sequence data will likely maximize the value of the data to the community (Field et al., 2009; Kaye et al., 2009; Whitlock et al., 2010), and encourage comparative studies as seen in other model organism communities such as ZFIN (Howe et al., 2012), WORM-BOOK (Greenwald, 2016), and FLYBASE (Attrill et al., 2016).

With an annotated genome and multiple assembled and annotated transcriptomes now available for electric fishes (and more currently being assembled – Gallant; unpublished data), the computational development of markers for future phylogenetic studies should be much more efficient. A collection of electric fish nucleotide and protein databases are available at <http://efishgenomics.integrativebiology.msu.edu/> and submissions of sequence data are welcomed (visit the site for details).

To maximize the value of resources, researchers need an efficient way to share genomic tools and established protocols. Like the aforementioned databases for other organisms, a putative 'EFISHBASE' should include databases and repositories for transgenic lines, CRISPR and RNAi components, plasmids, etc. with protocols for their use. In addition, tools to facilitate neuroscience studies (e.g., species brain atlases; video protocols for performing

neuronal recordings) and comparative work (e.g., tools for assessing gene homology, bioinformatics pipelines; sets of genetic markers) should be included along with the sequence databases currently available (marker datasets are available for most Mormyrid genera, but are not yet curated in a central repository).

3. Prospects for electric fish research – the next 30 years

Electric fish are a system with the potential to address many broad biological questions. Decades of research with traditional molecular tools on a variety of topics (Section 1) have been very productive; however, the development of molecular tools over the next decade or so (Section 2) will allow electric fish researchers to rapidly increase the pace of discovery. Applications of the genomic tools outlined above promise advances in electric fish research, and the ability to tackle some long-standing questions that have proven difficult to approach with pre-genomic tools. This section discusses a partial list of research avenues that we expect modern genomic tools to open up in the coming decades.

3.1. Gene expression profiles

First-generation sequencing approaches (Sanger sequencing of PCR products) to gene expression are usually limited to the study of one or a few genes simultaneously. Although valuable advances have been made with these techniques (Section 1), constructing a gene regulatory network using these methods is not practical. RNAseq can capture the entire expression profile of the target cells (Section 2.2), making it a formidable tool to describe gene networks and understand their consequences for tissue functions/development.

This strategy could determine whether compartmentalization of *scn4aa* and *scn4ab* expression occurs between subsets of muscle fibers in non-electrogenic fish (Zakon et al., 2008), and which gene interactions mediate such specialization. Likewise, single-cell RNAseq (Kolodziejczyk et al., 2015) could identify candidate gene(s) responsible for signal transduction after comparing activated and non-activated electroreceptor cells. Comparative expression studies could also elucidate the molecular specializations in other electric fish tissues; for example, those between motoneurons and electromotoneurons.

Tissue comparisons have proven useful for understanding the differentiation between EO and skeletal muscle. Lamanna et al. (2014) found very similar gene expression profiles between muscle and EO transcriptomes from *C. compressirostris*. Further work has discovered hundreds of differentially expressed genes between these tissues, and found that the EO exhibits higher gene expression variability than muscle, across *C. compressirostris*, *C. tshokwe*, and *Gnathonemus petersii* (Lamanna et al., 2015). These comparisons have also identified potential genes involved in specialized EO functions, such as cell size, structure and insulation (Gallant et al., 2014), ion pumps, and membrane synthesis and turnover (Lamanna et al., 2015).

Complementing RNAseq of mRNA with miRNA-sequencing has broadened understanding of how gene expression profiles are regulated in the EO. An EO-specific miRNA (Traeger et al., 2015) and three muscle-expressed miRNA are upregulated in the EO compared to muscle (Pinch et al., 2016). To our knowledge, ribosome profiling (Ribo-seq) has yet to be used in electric fish research, but it could potentially provide information about post-transcriptional regulation. Ribo-seq captures the subset of mRNAs that are translated at a particular moment (Ingolia et al., 2009), and thus can be used to study the translational control of gene expression (Ingolia, 2014). Further application of these technologies should provide valuable insights to understand the function, evolu-

tion, and diversification of EOs and the nervous system control and perception of EODs.

3.2. Function of electrocyte proteins

The aforementioned work on the voltage-gated Na⁺ channel gene *scn4aa* (Section 1.2) describes an example of a protein adapted for electrocyte function. Yet, Na⁺ channels are not solely responsible for the propagation of action potentials, let alone the entire function of the electrocyte. There are likely additional examples of proteins co-opted for the electrocyte phenotype. Paralogs are prime candidates for such differential expression among tissues; nevertheless it would be most interesting to learn how cartilaginous electric fish (i.e., fishes that did not experience the teleost-specific whole genome duplication) became electrogenic.

Potential candidates of specialized electrocyte proteins could include those mentioned above (Section 3.1) as well as proteins involved in hormone/neurotransmitter reception (Sections 1.2 and 1.3 respectively). Perhaps the most intuitive candidates should be K⁺ channels. Rashid and Dunn (1998) identified 19 potential K⁺ channel gene fragments in *Apteronotus leptorhynchus*, and suggested that K⁺ channels play critical roles in the precise regulation of neuronal firing that is required in complex nervous systems. Furthermore, rectifying K⁺ currents contribute to EOD production: K⁺ and Na⁺ kinetics are highly correlated in *Sternopygus*, reflecting a high degree of coregulation between two distinct ion channels (McAnelly and Zakon, 2000), yet the expression K⁺ and Na⁺ channels is extremely tissue-specific in *Eigenmannia virescens* (Ban et al., 2015). Therefore, it is worth exploring whether K⁺ channels have evolved in parallel to Na⁺ channels in Gymnotiform and Mormyroid electric fishes (Zakon et al., 2008). Whole-genome sequencing (see Section 2.2) could provide a comprehensive picture of the genes present and their homologies, suggest hypotheses about their evolutionary histories, and identify positively selected substitutions. The availability of well annotated genomes and transcriptomes would open the door to proteomic approaches; providing an opportunity to validate hypotheses in post-transcriptional regulation and alternative splicing. Additionally, predictions about proteins can be made from well annotated genomes/transcriptomes, allowing *in silico* trypsin-digested protein patterns to be generated as well as helping to identify proteins via mass spectrometry. The functional consequences of substitutions, alternative-splicing and post-transcriptional regulation could be inspected through the tools described in Sections 2.3 and 2.4.

3.3. Understanding the evolution of biodiversity using weakly electric fish as a model

With more NGS sequencing, electric fish may enable us to address some of the large-scale questions in evolutionary biology. Mormyrid and Gymnotiform weakly electric fish are remarkably speciose (Albert et al., 2005; Sullivan et al., 2000); their independent origins constitute a 'natural experiment' in species adaptation and radiation. In some electric fish clades, the changes involved in macroevolutionary diversification are mirrored at lower taxonomic levels; for example the repeated switch between bi- and tri-phasic EODs that seems to have occurred among genera and among conspecific populations in mormyrids (Arnegard and Carlson, 2005; Gallant et al., 2011). This is a truly rare opportunity to investigate potentially incipient speciation processes, with the aid of behavioral, life-history, and comparative genomics studies (Section 2.2).

Electric fish studies could address the relative importance – and potential conflict – between natural and sexual selection during the process of adaptive radiation (Andersson, 1994; Coyne and Orr, 2004; Schluter, 2001; Wagner et al., 2012). The EOD is used for species identification (Curtis and Stoddard, 2003; Feulner

et al., 2009; Kramer, 1997; Silva et al., 2008) and therefore EOD divergence may be implicated in the radiation of these groups through sexual selection. Detailed morphological, physiological, and behavioral studies on electric fishes have given us an enviable understanding of their communication systems (Baker et al., 2013; Carlson, 2002; Carlson and Gallant, 2013). Two integrative studies suggest that adaptation to local ecological conditions drive divergence among *Campylomormyrus* species (Feulner et al., 2008); whereas selection on sexual signals seems more important among the *Paramormyrops* (Arnegard et al., 2010). With genomic sequencing (Section 2.2), it will be possible to use population genetic analyses to identify regions of the genome that indicate positive selection; i.e., genes/functions that are driving species divergence (Zakon et al., 2006). Comparisons of these regions between different species/populations of electric fish would be a valuable addition to the understanding of population genetics.

3.4. Understanding the role of cis-regulatory changes in the evolution of development

The availability of genomic-level data has opened up a number of broad-scale questions in evolution and development, in particular; the relative contributions of change in protein-coding sequence vs. cis-regulatory elements (Hoekstra and Coyne, 2007). Whole genome sequencing holds the key for unlocking the role cis-regulatory elements may play in the evolution, differentiation and maintenance of EOs (Section 2.2).

Zakon et al. (2008) suggest that the comparison of transcription factor binding sites from the gene *scn4aa* between electric and non-electrogenic taxa could help explain the genetic underpinnings of *scn4aa* specialization to electrocytes. We propose expanding on this with comparative genomics between fully sequenced genomes. RNAseq (Section 2.1) could tackle this question much more broadly, by comparing additional regulatory elements, mechanisms, and genes involved in EO differentiation. We expect that a number of key signaling and regulatory molecules in electrocytes would have also evolved in concert with *scn4aa* and EOD signal divergence (Markham, 2013).

Once differences in cis-regulatory regions between genes/populations/species are identified, CRISPR (Section 2.4) can be utilized to assess their function. CRISPR can be directed to cis-regulatory regions, either to disrupt or replace the sequence (e.g., use HDR to replace an enhancer region for one gene with that of another gene, or of the same gene from a different species). CRISPR can be used to replace/alter transcription factor coding sequences, providing an opportunity to utilize electric fish as a model to address the relative impact of cis-regulation and coding sequence change in evolution (Hoekstra and Coyne, 2007; Schmidt et al., 2010).

3.5. Manipulation of neural circuits using transgenics

Numerous studies on electric fish have contributed to our current knowledge in systems and circuits neuroscience. Examples include: mechanisms for preservation and analysis of temporal information (Hopkins, 1999; Kawasaki, 1996), synaptic plasticity (Bell et al., 1997), cerebellar function (Sawtell and Bell, 2008), and effects of both motor (Bell and Grant, 1989) and neuromodulatory systems (Deemyad et al., 2013) on sensory processing. The neural circuitry underlying many complex electric fish behaviors has been well studied; the jamming avoidance response is completely understood from peripheral signal perception through motor output (Heiligenberg, 1991).

The electrosensory lobe is an important locus for understanding how neurons code information, i.e., how sensory perceptions are represented in configurations of all-or-none action potentials (Chacron et al., 2011; Clarke et al., 2015). Moreover, the repeated

independent evolution of electrosensory systems in electric fishes allows for the rare opportunity of comparing convergent electrosensory adaptations to common pressures related to information processing or behavioral control (Kawasaki, 1996).

Further progress could be fueled by using advanced neuroscientific tools, such as the development of transgenic electric fish lines for optogenetics. This technique involves the expression of light-sensitive ion channel proteins (e.g., channelrhodopsin or halorhodopsin) in particular neuronal populations, and allows researchers to manipulate neuronal activity with light (Deisseroth, 2010; Mancuso et al., 2010). Optogenetics allows for monitoring and manipulation of neural activity from intact animals, in real time, with minimal interference of behavior. Optogenetics could also provide an alternate method (vs. voltage- or patch-clamp recordings) for recording ionic currents in EOs and/or the brain in species/situations where electrophysiology may be difficult.

Use of an inducible construct (e.g., tamoxifen-induced) to over-express Caspase9 in specific neurons could facilitate studies of neuronal circuits. CRISPR could be used to direct, and insert, the inducible construct to the genome. Once integrated and induced, the overexpression of Caspase9 will lead to cell death via apoptosis (Druskovic et al., 2006), causing neuronal ablations. The comparison of pre- and post-treatment measurements could hint at the function of the lost brain region, as well as its role in behavior.

CRISPR can be used to generate 'multi-transgenic' fishes. For example, instead of tagging a single protein with GFP (see Section 2.4), other colored fluorescent proteins (e.g., RFP, YFP, BFP – see Dean and Palmer, 2014) can tag additional gene products, providing simultaneous comparisons of multiple proteins within a single fish (e.g., visualization of multiple neurotransmitter/hormones and their receptors in a single neuron, identification of neuronal subtypes/populations in whole brain sections). As mentioned in Section 2.4, *in situ* hybridization or fluorescent *in situ* hybridization in a 'multi-transgenic' fish can be used to label additional genes of interest that the transgenic fish doesn't express.

Once genome manipulation is established in electric fish (Section 2.4), there are further transgenic manipulations that could be applied to answer precise questions. One such technique is 'Brainbow' labeling where (potentially) each cell in an organism displays a different fluorescent profile that is unchanged after cell division, during morphogenesis, and during/after development (Pan et al., 2013). CRISPR can be used to incorporate the 'brainbow' construct into the genome through HDR (Section 2.4). This technique has been used in skin to monitor regeneration (Chen et al., 2016), and in neurons to trace brain connections (Pan et al., 2011). The possibility to study neuronal connections in electric fish using this technique is very alluring. Many of these proposed questions/techniques are within the first accessible applications of this research.

Conclusion – looking to the future

The weakly electric fish community has a strongly integrated concept of phenotype (Fig. 1), which stems from extensive characterization of natural behavior (because of its historical roots in neuroethology). Both biologists and the groups of fishes that they study are diverse; the weakly electric fish community is rich with mechanistic understanding of phenotypes over multiple levels of biological analysis. Because of the new arrival of low cost, high throughput sequencing technologies, these rich sources of phenotypic data position the electric fish model as a unique system to tackle highly integrative 21st century biology research objectives, such as understanding the brain and integrating phenotype and genotype. A concerted effort by researchers in the electric fish community focused on strategic sequencing and techniques development over the next decade could reap outsized rewards, and

position the community as a leader in bringing genomics techniques to non-model systems.

Acknowledgements

The authors wish to thanks K. Dunlap, A. Charbonneau, D. Tack, A. Parigi and K. Allen for their helpful comments on an earlier version of this manuscript. The authors would additionally like to acknowledge the organizers of the 2016 Weakly Electric Fish Meeting in Montevideo, Uruguay, as well as its many participants, for ideas and feedback that formed the basis of this review. The authors are supported by NSF Grant 1455405 to JRG.

References

- Alagia, A., Eritja, R., 2016. siRNA and RNAi optimization. *WIREs RNA* 7, 316–329.
- Albert, J.S., Crampton, W.G.R., Thorsen, D.H., Lovejoy, N.R., 2005. Phylogenetic systematics and historical biogeography of the Neotropical electric fish *Gymnotus* (Teleostei: Gymnotidae). *Syst. Biodivers.* 2, 375–417.
- Allee, S.J., Markham, M.R., Salazar, V.L., Stoddard, P.K., 2008. Opposing actions of 5HT1A and 5HT2-like serotonin receptors on modulations of the electric signal waveform in the electric fish *Brachyhyppopomus pinnicaudatus*. *Horm. Behav.* 53, 481–488.
- Allee, S.J., Markham, M.R., Stoddard, P.K., 2009. Androgens enhance plasticity of an electric communication signal in female knifefish, *Brachyhyppopomus pinnicaudatus*. *Horm. Behav.* 56, 264–273.
- Alves-Gomes, J., Hopkins, C.D., 1997. Molecular insights into the phylogeny of Mormyrid fishes and the evolution of their electric organs. *Brain Behav. Evol.* 49, 324–350.
- Amores, A., Force, A., Yan, Y.L., Joly, L., Amemiya, C., Fritz, A., Ho, R.K., Langeland, J., Prince, V., Wang, Y.L., Westerfield, M., Ekker, M., Postlethwait, J.H., 1998. Zebrafish hox clusters and vertebrate genome evolution. *Science* 282, 1711–1714.
- Andersson, M.B., 1994. Sexual Selection. Princeton Univ Pr.
- Andrews, O.E., Cha, D.J., Wei, C., Patton, J.G., 2014. RNAi-mediated gene silencing in zebrafish triggered by convergent transcription. *Sci. Rep.* 4, 1–8.
- Arnegard, M.E., Bogdanowicz, S.M., Hopkins, C.D., 2005. Multiple cases of striking genetic similarity between alternate electric fish signal morphs in sympatry. *Evolution* 59, 324.
- Arnegard, M.E., Carlson, B.A., 2005. Electric organ discharge patterns during group hunting by a mormyrid fish. *Proc. Biol. Sci.* 272, 1305–1314.
- Arnegard, M.E., Zwickl, D.J., Lu, Y., Zakon, H.H., Robinson, G.E., 2010. Old gene duplication facilitates origin and diversification of an innovative communication system—twice. *Proc. Natl. Acad. Sci. U.S.A.* 107, 22172–22177.
- Attrill, H., Falls, K., Goodman, J.L., Millburn, G.H., Antonazzo, G., Rey, A.J., Marygold, S.J., The FlyBase Consortium, 2016. FlyBase: establishing a Gene Group resource for *Drosophila melanogaster*. *Nucleic Acids Res.* 44, D786–D792.
- Auer, T.O., Durore, K., De Cian, A., Concordet, J.P., Del Bene, F., 2014. Highly efficient CRISPR/Cas9-mediated knock-in in zebrafish by homology-independent DNA repair. *Genome Res.* 24, 142–153.
- Baker, C.V.H., Modrell, M.S., Gillis, J.A., 2013. The evolution and development of vertebrate lateral line electroreceptors. *J. Exp. Biol.* 216, 2515–2522.
- Ban, Y., Smith, B.E., Markham, M.R., 2015. A highly polarized excitable cell separates sodium channels from sodium-activated potassium channels by more than a millimeter. *J. Neurophysiol.* 114, 520–530.
- Barrangou, R., Fremaux, C., Deveau, H., Richards, M., Boyaval, P., Moineau, S., Romero, D.A., Horvath, P., 2007. CRISPR provides acquired resistance against viruses in prokaryotes. *Science* 315, 1709–1712.
- Bass, A.H., Hopkins, C.D., 1985. Hormonal control of sex differences in the electric organ discharge (EOD) of mormyrid fishes. *J. Comp. Physiol. A* 156, 587–604.
- Bass, A.H., Hopkins, C.D., 1984. Shifts in frequency tuning of electroreceptors in androgen-treated mormyrid fish. *J. Comp. Physiol.* 155, 713–724.
- Bastian, J., Schniederjan, S., Nguyenkim, J., 2001. Arginine vasotocin modulates a sexually dimorphic communication behavior in the weakly electric fish *Apteronotus leptorhynchus*. *J. Exp. Biol.* 204, 1909–1924.
- Bell, C.C., Grant, K., 1989. Corollary discharge inhibition and preservation of temporal information in a sensory nucleus of mormyrid electric fish. *J. Neurosci.* 9 (3), 1029–1044.
- Bell, C.C., Han, V.Z., Sugawara, Y., Grant, K., 1997. Synaptic plasticity in a cerebellum-like structure depends on temporal order. *Nature* 387, 278–281.
- Bensouilah, M., Schugardt, C., Roesler, R., Kirschbaum, F., Denizot, J.-P., 2002. Larval electroreceptors in the epidermis of mormyrid fish: I. Tuberos organs of type A and B. *J. Comp. Neurol.* 447, 309–322.
- Bild, A.H., Chang, J.T., Johnson, W.E., Piccolo, S.R., 2014. A field guide to genomics research. *PLoS Biol.* 12, e1001744.
- Borgman, C.L., 2012. The conundrum of sharing research data. *Acta Anaesthesiol. Scand.* 63, 1059–1078.
- Bottai, D., Dunn, R.J., Ellis, W., Maler, L., 1997. N-methyl-D-aspartate receptor 1 mRNA distribution in the central nervous system of the weakly electric fish *Apteronotus leptorhynchus*. *J. Comp. Neurol.* 389, 65–80.
- Bradnam, K.R., Fass, J.N., Alexandrov, A., Baranay, P., Bechner, M., Birol, I., Boisvert, S., Chapman, J.A., Chapuis, G., Chikhi, R., Chitsaz, H., Chou, W.-C., Corbeil, J., Del Fabbro, C., Docking, T., Durbin, R., Earl, D., Emrich, S., Fedotov, P., Fonseca, N.A., Ganapathy, G., Gibbs, R.A., Gnerre, S., Godzaridis, E., Goldstein, S., Haimel, M., Hall, G., Haussler, D., Hiatt, J.B., Ho, I.Y., Howard, J., Hunt, M., Jackman, S.D., Jaffe, D.B., Jarvis, E.D., Jiang, H., Kazakov, S., Kersey, P.J., Kitzman, J.O., Knight, J.R., Koren, S., Lam, T.-W., Lavenier, D., Laviolette, F., Li, Y., Li, Z., Liu, B., Liu, Y., Luo, R., MacCallum, I., MacManes, M.D., Maillet, N., Melnikov, S., Naquin, D., Ning, Z., Otto, T.D., Paten, B., Paulo, O.S., Philipp, A.M., Pina-Martins, F., Place, M., Przybylski, D., Qin, X., Qu, C., Ribeiro, F.J., Richards, S., Rokhsar, D.S., Ruby, J., Scalabrin, S., Schatz, M.C., Schwartz, D.C., Sergushichev, A., Sharpe, T., Shaw, T.I., Shendure, J., Shi, Y., Simpson, J.T., Song, H., Tsarev, F., Vezzi, F., Vicedomini, R., Vieira, B.M., Wang, J., Worley, K.C., Yin, S., Yiu, S.-M., Yuan, J., Zhang, G., Zhang, H., Zhou, S., Korf, I.F., 2013. Assemblathon 2: evaluating de novo methods of genome assembly in three vertebrate species. *GigaScience* 2, 10–31.
- Buffalo, V., 2015. Bioinformatics Data Skills. O'Reilly Media.
- Carlson, B.A., 2002. Neuroanatomy of the mormyrid electromotor control system. *J. Comp. Neurol.* 454, 440–455.
- Carlson, B.A., Gallant, J.R., 2013. From sequence to spike to spark: Evo-devo-neuroethology of electric communication in mormyrid fishes. *J. Neurogenet.* 27, 106–129.
- Carlson, B.A., Hasan, S.M., Hollmann, M., Miller, D.B., Harmon, L.J., Arnegard, M.E., 2011. Brain evolution triggers increased diversification of electric fishes. *Science* 332, 583–586.
- Carlson, B.A., Hopkins, C.D., Thomas, P., 2000. Androgen correlates of socially induced changes in the electric organ discharge waveform of a mormyrid fish. *Horm. Behav.* 38, 177–186.
- Chacron, M.J., Longtin, A., Maler, L., 2011. Efficient computation via sparse coding in electrosensory neural networks. *Curr. Opin. Neurobiol.* 21, 752–760.
- Changeux, J.P., 2012. The nicotinic acetylcholine receptor: the founding father of the pentameric ligand-gated ion channel superfamily. *J. Biol. Chem.* 287, 40207–40215.
- Changeux, J.P., Kasai, M., Lee, C.Y., 1970. Use of a snake venom toxin to characterize the cholinergic receptor protein. *Proc. Natl. Acad. Sci.* 67, 1241–1247.
- Chen, C.-H., Puliafito, A., Cox, B.D., Primo, L., Fang, Y., Di Talia, S., Poss, K.D., 2016. Multicolor cell barcoding technology for long-term surveillance of epithelial regeneration in zebrafish. *Dev. Cell* 36, 668–680.
- Clarke, S.E., Longtin, A., Maler, L., 2015. The neural dynamics of sensory focus. *Nat. Commun.* 6, 8764.
- Coyne, J.A., Orr, H.A., 2004. Speciation. Sinauer Associates Incorporated.
- Cuellar, H., Kim, J.A., Unguez, G.A., 2006. Evidence of post-transcriptional regulation in the maintenance of a partial muscle phenotype by electrogenic cells of *S. macrurus*. *FASEB J.* 20, E1856–E1865.
- Curtis, C.C., Stoddard, P.K., 2003. Mate preference in female electric fish, *Brachyhyppopomus pinnicaudatus*. *Anim. Behav.* 66, 329–336.
- Davey, J.W., Blaxter, M.L., 2011. RADSeq: next-generation population genetics. *Brief. Funct. Genom.* 9, 416–423.
- Dean, K.M., Palmer, A.E., 2014. Advances in fluorescence labeling strategies for dynamic cellular imaging. *Nat. Chem. Biol.* 10 (7), 512–523.
- De Rienzo, G., Gutzman, J.H., Sive, H., 2012. Efficient shRNA-mediated inhibition of gene expression in zebrafish. *Zebrafish* 9, 97–107.
- De Wit, P., Pespeni, M.H., Palumbi, S.R., 2015. SNP genotyping and population genomics from expressed sequences – current advances and future possibilities. *Mol. Ecol.* 24, 2310–2323.
- Deemyad, T., Metzner, M.G., Pan, Y., 2013. Serotonin selectively enhances perception and sensory neural responses to stimuli generated by same-sex conspecifics. *Proc. Natl. Acad. Sci. U.S.A.* 110 (48), 19609–19614.
- Deisseroth, K., 2010. Optogenetics. *Nat. Methods* 8, 26–29.
- Deisseroth, K., Feng, G., Majewska, A.K., Miesenböck, G., Ting, A., Schnitzer, M.J., 2006. Next-generation optical technologies for illuminating genetically targeted brain circuits. *J. Neurosci.* 26, 10380–10386.
- Denizot, J.-P., Bensouilah, M., Roesler, R., Schugardt, C., Kirschbaum, F., 2007. Larval electroreceptors in the epidermis of mormyrid fish: II. The promormyromast. *J. Comp. Neurol.* 501, 810–823.
- Diedhiou, S., Bartsch, P., Kirschbaum, F., 2007. The embryonic and larval development of *Pollimyrus isidori* (Mormyridae, Osteoglossomorpha): its staging with reference to structure and behavior. *Bull. Fish Biol.* 9, 61–88.
- Doench, J.G., Hartenian, E., Graham, D.B., Tothova, Z., Hegde, M., Smith, I., Sullender, M., Ebert, B.L., Xavier, R.J., Root, D.E., 2014. Rational design of highly active sgRNAs for CRISPR-Cas9-mediated gene inactivation. *Nat. Biotechnol.* 32, 1262–1267.
- Druskovic, M., Suput, D., Milisav, I., 2006. Overexpression of caspase-9 triggers its activation and apoptosis *in vitro*. *Croat. Med. J.* 47, 832–840.
- Dulka, J.G., Maler, L., Ellis, W., 1995. Androgen-induced changes in electrocommunication behavior are correlated with changes in substance P-like immunoreactivity in the brain of the electric fish *Apteronotus leptorhynchus*. *J. Neurosci.* 15, 1879–1890.
- Dunlap, K.D., Castellano, J.F., Prendaj, E., 2006. Social interaction and cortisol treatment increase cell addition and radial glia fiber density in the diencephalic periventricular zone of adult electric fish, *Apteronotus leptorhynchus*. *Horm. Behav.* 50, 10–17.
- Dunlap, K.D., McAnelly, M.L., Zakon, H.H., 1997. Estrogen modifies an electrocommunication signal by altering the electrocyte sodium current in an electric fish, *Sternopygus*. *J. Neurosci.* 17, 2869–2875.
- Dunlap, K.D., Zakon, H.H., 1998. Behavioral actions of androgens and androgen receptor expression in the electrocommunication system of an electric fish, *Eigenmannia virescens*. *Horm. Behav.* 34, 30–38.

- Kotb, A.M., Simon, O., Blumenthal, A., Vogelgesang, S., Dombrowski, F., Amann, K., Zimmermann, U., Endlich, K., Endlich, N., 2016. Knockdown of ApoL1 in zebrafish larvae affects the glomerular filtration barrier and the expression of nephrin. *PLoS ONE* 11, e0153768–17.
- Kramer, B., 1997. Electric organ discharges and their relation to sex in mormyrid fishes. *Naturwissenschaften* 84, 119–121.
- Lamanna, F., Kirschbaum, F., Tiedemann, R., 2014. De novo assembly and characterization of the skeletal muscle and electric organ transcriptomes of the African weakly electric fish *Campylomormyrus compressirostris* (Mormyridae, Teleostei). *Mol. Ecol. Resour.* 14, 1222–1230.
- Lamanna, F., Kirschbaum, F., Waurick, I., Dieterich, C., Tiedemann, R., 2015. Cross-tissue and cross-species analysis of gene expression in skeletal muscle and electric organ of African weakly-electric fish (Teleostei; Mormyridae). *BMC Genomics* 16, 1–17.
- Lavoué, S., Miya, M., Arnegard, M.E., Sullivan, J.P., Hopkins, C.D., Nishida, M., 2012. Comparable ages for the independent origins of electrogenesis in African and South American weakly electric fishes. *PLoS ONE* 7, e36287–18.
- Lavoué, S., Sullivan, J.P., Arnegard, M.E., Hopkins, C.D., 2008. Differentiation of morphology, genetics and electric signals in a region of sympatry between sister species of African electric fish (Mormyridae). *J. Evol. Biol.* 21, 1030–1045.
- Lee, H., Gurtowski, J., Yoo, S., Nattestad, M., Marcus, S., 2016. Third-generation sequencing and the future of genomics. *bioRxiv*.
- Li, J., Zhang, B.-B., Ren, Y.-G., Gu, S.-Y., Xiang, Y.-H., Du, J.-L., 2015. Intron targeting-mediated and Endogenous Gene Integrity-maintaining Knockin in Zebrafish using the CRISPR/Cas9 System, vol. 25. Nature Publishing Group, pp. 634–637.
- Li, Q., Lee, J.-A., Black, D.L., 2007. Neuronal regulation of alternative pre-mRNA splicing. *Nat. Rev. Neurosci.* 8, 819–831.
- Lissmann, H.W., 1958. On the function and evolution of electric organs in fish. *J. Exp. Biol.* 35, 156–191.
- Liu, E.T., 2005. Systems biology, integrative biology, predictive biology. *Cell* 121, 505–506.
- Lohman, B.K., Weber, J.N., Bolnick, D.I., 2016. Evaluation of TagSeq, a reliable low-cost alternative for RNAseq. *Mol. Ecol. Resour.* 16, 1315–1321.
- Lopreato, G.F., Lu, Y., Southwell, A., Atkinson, N.S., Hillis, D.M., Wilcox, T.P., Zakon, H. H., 2001. Evolution and divergence of sodium channel genes in vertebrates. *Proc. Natl. Acad. Sci.* 98, 7588–7592.
- Macaulay, I.C., Voet, T., 2014. Single cell genomics: advances and future perspectives. *PLoS Genet.* 10, e1004126–9.
- Magadum, S., Banerjee, U., Murugan, P., Gangapur, D., Ravikesavan, R., 2013. Gene duplication as a major force in evolution. *J. Genet.* 92, 155–161.
- Maler, L., Ellis, W.G., 1987. Inter-male aggressive signals in weakly electric fish are modulated by monoamines. *Behav. Brain Res.* 25, 75–81.
- Maler, L., Monaghan, D., 1991. The distribution of excitatory amino acid binding sites in the brain of an electric fish, *Apteronotus leptorhynchus*. *J. Chem. Neuroanat.* 4, 39–61.
- Mancuso, J.J., Kim, J., Lee, S., Tsuda, S., Chow, N.B.H., Augustine, G.J., 2010. Optogenetic probing of functional brain circuitry. *Exp. Physiol.* 96, 26–33.
- Markham, M.R., 2013. Electrocyte physiology: 50 years later. *J. Exp. Biol.* 216, 2451–2458.
- Markham, M.R., Allee, S.J., Goldina, A., Stoddard, P.K., 2009a. Melanocortins regulate the electric waveforms of gymnotiform electric fish. *Horm. Behav.* 55, 306–313.
- Markham, M.R., McAnelly, M.L., Stoddard, P.K., Zakon, H.H., 2009b. Circadian and social cues regulate ion channel trafficking. *PLoS Biol.* 7, e1000203–14.
- Markham, M.R., Stoddard, P.K., 2013. Cellular mechanisms of developmental and sex differences in the rapid hormonal modulation of a social communication signal. *Horm. Behav.* 63, 586–597.
- Markham, M.R., Stoddard, P.K., 2005. Adrenocorticotrophic hormone enhances the masculinity of an electric communication signal by modulating the waveform and timing of action potentials within individual cells. *J. Neurosci.* 25, 8746–8754.
- Martin, J.A., Wang, Z., 2011. Next-Generation Transcriptome Assembly, vol. 12. Nature Publishing Group, pp. 671–682.
- Martin, S.H., Jiggins, C.D., 2013. Genomic studies of adaptation in natural populations. *eLS*.
- Maxwell, E.E., Fröbisch, N.B., Heppleston, A.C., 2008. Variability and conservation in Late Chondrichthyan Development: ontogeny of the Winter Skate (*Leucoraja ocellata*). *Anat. Rec.* 291, 1079–1087.
- McAnelly, L., Silva, A., Zakon, H.H., 2003. Cyclic AMP modulates electrical signaling in a weakly electric fish. *J. Comp. Physiol. A* 189, 273–282.
- McAnelly, L., Zakon, H.H., 1996. Protein kinase A activation increases sodium current magnitude in the electric organ of *Sternopygus*. *J. Neurosci.* 16, 4383–4388.
- McAnelly, M.L., Zakon, H.H., 2007. Androgen modulates the kinetics of the delayed rectifying K⁺ current in the electric organ of a weakly electric fish. *Dev. Neurobiol.* 67, 1589–1597.
- McAnelly, M.L., Zakon, H.H., 2000. Coregulation of voltage-dependent kinetics of Na⁺ and K⁺ currents in electric organ. *J. Neurosci.* 20, 3408–3414.
- Mei, Y., Wang, Y., Chen, H., Sun, Z.S., Ju, X.-D., 2016. Recent progress in CRISPR/Cas9 technology. *J. Genet. Genom.* 43, 63–75.
- Metzker, M.L., 2009. Sequencing Technologies – The Next Generation, vol. 11. Nature Publishing Group, pp. 31–46.
- Miledi, R., Molinoff, P., Potter, L.T., 1971. Biological sciences: isolation of the cholinergic receptor protein of torpedo electric tissue. *Nature* 229, 554–557.
- Miller, C., 2000. Ion channel surprises. *Neuron* 25, 7–9.
- Mills, A., Zakon, H.H., 1987. Coordination of Eod frequency and pulse duration in a weakly electric wave fish – the influence of androgens. *J. Comp. Physiol. A* 161, 417–430.
- Mitra, A.K., McCarthy, M.P., Stroud, R.M., 1989. Three-dimensional structure of the nicotinic acetylcholine receptor and location of the major associated 43-kD cytoskeletal protein, determined at 22 Å by low dose electron microscopy and x-ray diffraction to 12.5 Å. *J. Cell Biol.* 109, 755–774.
- Morcos, P.A., Vincent, A.C., Moulton, J.D., 2015. Gene editing versus morphants. *Zebrafish* 12, 319.
- Mortazavi, A., Williams, B.A., McCue, K., Schaeffer, L., Wold, B., 2008. Mapping and quantifying mammalian transcriptomes by RNA-Seq. *Nat. Methods* 5, 621–628.
- Mou, T., Kraas, J.R., Fung, E.T., Swope, S.L., 1998. Identification of a dynein molecular motor component in Torpedo electroplax; binding and phosphorylation of Tctex-1 by Fyn. *FEBS Lett.* 435, 275–281.
- Moulton, J.D., 2016. Guide for morpholino users: Toward therapeutics. *J. Drug Discov. Develop. Deliv.* 3, 270–282.
- Nagamachi, C.Y., Pieczarka, J.C., Milhomem, S.S.R., O'Brien, P.C.M., de Souza, A.C.P., Ferguson-Smith, M.A., 2010. Multiple rearrangements in cryptic species of electric knifefish, *Gymnotus carapo* (Gymnotidae, Gymnotiformes) revealed by chromosome painting. *BMC Genet.* 11, 28.
- Narum, S.R., Buerkle, C.A., Davey, J.W., Miller, M.R., Hohenlohe, P.A., 2013. Genotyping-by-sequencing in ecological and conservation genomics. *Mol. Ecol.* 22, 2841–2847.
- National Research Council Committee on Research at the Intersection of the Physical and Life Sciences, Board on Physics and Astronomy, Board on Life Sciences, Board on Chemical Sciences and Technology, Division on Engineering and Physical Sciences, Division on Earth and Life Studies, 2010. Research at the Intersection of the Physical and Life Sciences. National Academies Press.
- Nazarian, J., Berry, D.L., Sanjari, S., Razvi, M., Brown, K., Hathout, Y., Vertes, A., Dadgar, S., Hoffman, E.P., 2011. Evolution and comparative genomics of subcellular specializations: EST sequencing of *Torpedo* electric organ. *Mar. Genom.* 4, 33–40.
- Nazarian, J., Hathout, Y., Vertes, A., Hoffman, E.P., 2007. The proteome survey of an electricity-generating organ (*Torpedo californica* electric organ). *Proteomics* 7, 617–627.
- Nitkin, R.M., Smith, M.A., Magill, C., Fallon, J.R., Yao, Y.M., Wallace, B.G., McMahan, U.J., 1987. Identification of agrin, a synaptic organizing protein from *Torpedo* electric organ. *J. Cell Biol.* 105, 2471–2478.
- Notch, E.G., Shaw, J.R., Coutermarsh, B.A., Dzioba, M., Stanton, B.A., 2011. Morpholino gene knockdown in adult *Fundulus heteroclitus*: Role of SGK1 in seawater acclimation. *PLoS ONE* 6, e29462–9.
- Novak, A.E., Jost, M.C., Lu, Y., Taylor, A.D., Zakon, H.H., Ribera, A.B., 2006. Gene duplications and evolution of vertebrate voltage-gated sodium channels. *J. Mol. Evol.* 63, 208–221.
- Nunes, M.D.S., Arif, S., Schlotterer, C., McGregor, A.P., 2013. A perspective on micro-Evo-Devo: progress and potential. *Genetics* 195, 625–634.
- Ohno, S., 2013. Evolution by Gene Duplication. Springer Science & Business Media, Berlin, Heidelberg.
- Paintner, S., Kramer, B., 2003. Electrosensory basis for individual recognition in a weakly electric, mormyrid fish, *Pollimyrus adspersus* (Günther, 1866). *Behav. Ecol. Sociobiol.* 55, 197–208.
- Pan, Y.A., Freundlich, T., Weissman, T.A., Schoppik, D., Wang, X.C., Zimmerman, S., Ciruna, B., Sanes, J.R., Lichtman, J.W., Schier, A.F., 2013. Zebrafish: multispectral cell labeling for cell tracing and lineage analysis in zebrafish. *Development* 140, 2835–2846.
- Pan, Y.A., Livet, J., Sanes, J.R., Lichtman, J.W., Schier, A.F., 2011. Multicolor Brainbow imaging in zebrafish. *Cold Spring Harb Protoc* 2011, pdb.prot5546.
- Patterson, J.M., Zakon, H.H., 1996. Differential expression of proteins in muscle and electric organ, a muscle derivative. *J. Comp. Neurol.* 370, 367–376.
- Patterson, J.M., Zakon, H.H., 1993. Bromodeoxyuridine labeling reveals a class of satellite-like cells within the electric organ. *J. Neurobiol.* 24, 660–674. <http://dx.doi.org/10.1002/neu.480240510>.
- Perrone, 2010. Vasotocin actions on electric behavior: interspecific, seasonal, and social context-dependent differences. *Front. Behav. Neurosci.* 1–12.
- Pinch, M., Güth, R., Samanta, M.P., Chaidze, A., Unguez, G.A., 2016. The myogenic electric organ of *Sternopygus macrurus*: a non-contractile tissue with a skeletal muscle transcriptome. *PeerJ* 4, e1828–30.
- Ran, F.A., Hsu, P.D., Lin, C.-Y., Gootenberg, J.S., Konermann, S., Trevino, A.E., Scott, D. A., Inoue, A., Matoba, S., Zhang, Y., Zhang, F., 2013. Double nicking by RNA-guided CRISPR Cas9 for enhanced genome editing specificity. *Cell* 154, 1380–1389.
- Rashid, A.J., Dunn, R.J., 1998. Sequence diversity of voltage-gated potassium channels in an electric fish. *Brain Res. Mol. Brain Res.* 54, 101–107.
- Rashid, A.J., Dunn, R.J., Turner, R.W., 2001. A prominent soma-dendritic distribution of Kv3.3 K⁺ channels in electrosensory and cerebellar neurons. *J. Comp. Neurol.* 441, 234–247.
- Rieger, F., Bon, S., Massoulié, J., Cartauld, J., Picard, B., Benda, P., 1976. *Torpedo marmorata* acetylcholinesterase: a comparison with the *Electrophorus electricus* enzyme. Molecular forms, subunits, electron microscopy, immunological relationship. *Eur. J. Biochem.* 68, 513–521.
- Salazar, V.L., Stoddard, P.K., 2009. Social competition affects electric signal plasticity and steroid levels in the gymnotiform fish *Brachyhypopomus gauderio*. *Horm. Behav.* 56, 399–409.
- Salisbury, J.P., Sîrbulescu, R.F., Moran, B.M., Auclair, J.R., Zupanc, G.K.H., Agar, J.N., 2015. The central nervous system transcriptome of the weakly electric brown

- ghost knifefish (*Apteronotus leptorhynchus*): de novo assembly, annotation, and proteomics validation. *BMC Genomics* 16, 166.
- Sander, J.D., Joung, J.K., 2014. CRISPR-Cas systems for editing, regulating and targeting genomes. *Nat. Biotechnol.* 32, 347–355.
- Sas, E., Maler, L., Tinner, B., 1990. Catecholaminergic systems in the brain of a gymnotiform teleost fish: an immunohistochemical study. *J. Comp. Neurol.* 292, 127–162.
- Sawtell, N.B., Bell, C.C., 2008. Adaptive processing in electrosensory systems: links to cerebellar plasticity and learning. *J. Physiol. Paris* 102, 223–232.
- Schluter, D., 2001. Ecology and the origin of species. *Trends Ecol. Evol.* 16, 372–380.
- Schmidt, D., Wilson, M.D., Ballester, B., Schwalie, P.C., Brown, G.D., Marshall, A., Kutter, C., Watt, S., Martinez-Jimenez, C.P., Mackay, S., Talianidis, I., Flicek, P., Odom, D.T., 2010. Five-vertebrate ChIP-seq reveals the evolutionary dynamics of transcription factor binding. *Science* 328, 1036–1040.
- Schugardt, C., Kirschbaum, F., 2004. Control of gonad maturation and regression by experimental variation of environmental factors in the mormyrid fish *Mormyrus rume probosciostris*. *Environ. Biol. Fishes* 70, 227–233.
- Schulman, S., Bendapudi, P., Sharda, A., Chen, V., Bellido-Martin, L., Jasuja, R., Furie, B.C., Flaumenhaft, R., Furie, B., 2016. Extracellular thiol isomerases and their role in thrombus formation. *Antioxid. Redox Signal.* 24, 1–15.
- Schwassmann, H.O., Assunção, M.I.S., Kirschbaum, F., 2014. Ontogeny of the electric organs in the electric eel, *Electrophorus electricus*: physiological, histological, and fine structural investigations. *Brain Behav. Evol.*, 1–16.
- Shestopalov, I.A., Pitt, C.L.W., Chen, J.K., 2012. Spatiotemporal resolution of the Ntla transcriptome in axial mesoderm development. *Nat. Chem. Biol.* 8, 270–276.
- Silva, A., Quintana, L., Perrone, R., Sierra, F., 2008. Sexual and seasonal plasticity in the emission of social electric signals. Behavioral approach and neural bases. *J. Physiol. Paris* 102, 272–278.
- Silva, D.S., Peixoto, L.A.W., Pieczarka, J.C., Wosiacki, W.B., Ready, J.S., Nagamachi, C. Y., 2015. Karyotypic and morphological divergence between two cryptic species of *Eigenmannia* in the Amazon basin with a new occurrence of XX/XY sex chromosomes (Gymnotiformes: Sternopygidae). *Neotrop. Ichthyol.* 13, 297–308.
- Sims, D., Sudbery, I., Illott, N.E., Heger, A., Ponting, C.P., 2014. Sequencing Depth and Coverage: Key Considerations in Genomic Analyses. *Nature Publishing Group* 15, 121–132.
- Smith, G.T., 2013. Evolution and hormonal regulation of sex differences in the electrocommunication behavior of ghost knifefishes (Apteronotidae). *J. Exp. Biol.* 216, 2421–2433.
- Smith, G.T., Combs, N., 2008. Serotonergic activation of 5HT1A and 5HT2 receptors modulates sexually dimorphic communication signals in the weakly electric fish *Apteronotus leptorhynchus*. *Horm. Behav.* 54, 69–82.
- Smith, G.T., Lu, Y., Zakon, H.H., 2000. Parvocells: a novel interneuron type in the pacemaker nucleus of a weakly electric fish. *J. Comp. Neurol.* 423, 427–439.
- Sobel, A., Weber, M., Changeux, J.P., 1977. Large-scale purification of the acetylcholine-receptor protein in its membrane-bound and detergent-extracted forms from *Torpedo marmorata* electric organ. *Eur. J. Biochem.* 80, 215–224.
- Stoddard, P.K., Markham, M.R., Salazar, V.L., Allee, S., 2007. Circadian rhythms in electric waveform structure and rate in the electric fish *Brachyhypopomus pinnicaudatus*. *Physiol. Behav.* 90, 11–20.
- Stroh, T., Zupanc, G.K.H., 1995. Somatostatin in the prepacemaker nucleus of weakly electric fish, *Apteronotus leptorhynchus*: evidence for a nonsynaptic function. *Brain Res.* 674, 1–14.
- Sullivan, J.P., Lavoué, S., Arnegard, M.E., Hopkins, C.D., 2004. AFLPs resolve phylogeny and reveal mitochondrial introgression within a species flock of African electric fish (Mormyroidea: Teleostei). *Evol.* 58, 825–838.
- Sullivan, J.P., Lavoué, S., Hopkins, C.D., 2000. Molecular systematics of the African electric fishes (Mormyroidea: teleostei) and a model for the evolution of their electric organs. *J. Exp. Biol.* 203, 665–683.
- Szabo, T., 1965. Sense organs of the lateral line system in some electric fish of the Gymnotidae, Mormyridae and Gymnarchidae. *J. Morphol.* 117, 229–249.
- Szabo, T., 1960. Development of the electric organ of Mormyridae. *Nature* 188, 760–762.
- Szabo, T., Kirschbaum, F., 1983. On the differentiation of electric organs in the absence of central connections or peripheral innervation. *The Physiology of Excitable Cells*, 451–460.
- Tang, X., Maegawa, S., Weinberg, E.S., Dmochowski, I.J., 2007. Regulating gene expression in zebrafish embryos using light-activated, negatively charged peptide nucleic acids. *J. Am. Chem. Soc.* 129, 11000–11001.
- Taylor, J.S., Van de Peer, Y., Braasch, I., Meyer, A., 2001. Comparative genomics provides evidence for an ancient genome duplication event in fish. *Philos. Trans. Roy. Soc. B: Biol. Sci.* 356, 1661–1679.
- Telgkamp, P., Combs, N., Smith, G.T., 2007. Serotonin in a diencephalic nucleus controlling communication in an electric fish: sexual dimorphism and relationship to indicators of dominance. *Dev. Neurobiol.* 67, 339–354.
- Terleph, T.A., 2003. Effects of social interaction on the electric organ discharge in a mormyrid fish, *Gnathonemus petersii* (Mormyridae, Teleostei). *J. Exp. Biol.* 206, 2355–2362.
- Thompson, A., Vo, D., Comfort, C., Zakon, H.H., 2014. Expression evolution facilitated the convergent neofunctionalization of a sodium channel gene. *Mol. Biol. Evol.* 31, 1941–1955.
- Traeger, L.L., Volkening, J.D., Moffett, H., Gallant, J.R., Chen, P.-H., Novina, C.D., Phillips, G.N., Anand, R., Wells, G.B., Pinch, M., Güth, R., Unguez, G.A., Albert, J.S., Zakon, H., Sussman, M.R., Samanta, M.P., 2015. Unique patterns of transcript and miRNA expression in the South American strong voltage electric eel (*Electrophorus electricus*). *BMC Genomics* 16, 243.
- Tsai, S.Q., Joung, J.K., 2016. Defining and improving the Genome-wide Specificities of CRISPR-Cas9 Nucleases, vol. 17. *Nature Publishing Group*, pp. 300–312.
- Unguez, G.A., Zakon, H.H., 2002. Skeletal muscle transformation into electric organ in *S. macrurus* depends on innervation. *J. Neurobiol.* 53, 391–402.
- Unguez, G.A., Zakon, H.H., 1998a. Reexpression of myogenic proteins in mature electric organ after removal of neural input. *J. Neurosci.* 18, 9924–9935.
- Unguez, G.A., Zakon, H.H., 1998b. Phenotypic conversion of distinct muscle fiber populations to electrocytes in a weakly electric fish. *J. Comp. Neurol.* 399, 20–34.
- VanGuilder, H., Vrana, K., Freeman, W., 2008. Twenty-five years of quantitative PCR for gene expression analysis. *Biotechniques* 44 (Supplement), 619–626.
- Wagner, C.E., Harmon, L.J., Seehausen, O., 2012. Ecological opportunity and sexual selection together predict adaptive radiation. *Nature*, 1–5.
- Wake, M.H., 2008. Integrative biology: science for the 21st century. *Bioscience* 58, 349–353.
- Weber, C.M., Martindale, M.Q., Tapscott, S.J., Unguez, G.A., 2012. Activation of Pax7-positive cells in a non-contractile tissue contributes to regeneration of myogenic tissues in the electric fish *S. macrurus*. *PLoS ONE* 7, e36819–14.
- Whitlock, M.C., McPeck, M.A., Rausher, M.D., Rieseberg, L., Moore, A.J., 2010. Data archiving. *Am. Nat.* 175, 145–146.
- Wong, T.-T., Zohar, Y., 2015. Production of reproductively sterile fish by a non-transgenic gene silencing technology. *Sci. Rep.*, 1–8.
- Yamamoto, T., Maler, L., Nagy, J.L., 1992. Organization of galanin-like immunoreactive neuronal systems in weakly electric fish (*Apteronotus leptorhynchus*). *J. Chem. Neuroanat.* 5, 19–38.
- Zakon, H.H., Lu, Y., Zwickl, D.J., Hillis, D.M., 2006. Sodium channel genes and the evolution of diversity in communication signals of electric fishes: convergent molecular evolution. *Proc. Natl. Acad. Sci.* 103, 3675–3680.
- Zakon, H.H., Unguez, G.A., 1999. Development and regeneration of the electric organ. *J. Exp. Biol.* 202, 1427–1434.
- Zakon, H.H., Zwickl, D.J., Lu, Y., Hillis, D.M., 2008. Molecular evolution of communication signals in electric fish. *J. Exp. Biol.* 211, 1814–1818.
- Zhang, J., 2003. Evolution by gene duplication: an update. *Trends Ecol. Evol.* 18, 292–298.
- Zhang, X.-H., Tee, L.Y., Wang, X.-G., Huang, Q.-S., Yang, S.-H., 2015. Off-target effects in CRISPR/Cas9-mediated genome. *Engineering*, 1–8.
- Zupanc, G.K., Okawara, Y., Zupanc, M.M., Fryer, J.N., Maler, L., 1991. *In situ* hybridization of putative somatostatin mRNA in the brain of electric gymnotiform fish. *NeuroReport* 2, 707–710.
- Zupanc, G.K.H., 2002. From oscillators to modulators: behavioral and neural control of modulations of the electric organ discharge in the gymnotiform fish, *Apteronotus leptorhynchus*. *J. Physiol. Paris* 96, 459–472.